arXiv:1508.00648v3 [math.CA] 11 Aug 2015

INTEGRAL REPRESENTATION OF WEIL'S ELLIPTIC FUNCTIONS

SU HU AND MIN-SOO KIM

Abstract. In this paper, we give a two dimensional analogue of the Euler-MacLaurin summation formula. By using this formula, we obtain an integral representation of Weil's elliptic functions which was introduced in the book "Elliptic functions according to Eisenstein and Kronecker".

1. Introduction

The Hurwitz-Lerch zeta function $\Phi(z,s,a)$ is defined by

$$\Phi(z,s,a)=\sum_{n=0}^{\infty}\frac{z^n}{(a+n)^s} \tag{1.1}$$

for

$$a\in\mathbb{C}\setminus\mathbb{Z}_0^-;\quad s\in\mathbb{C}\text{ when }|z|<1;\quad \Re(s)>k\ (k\in\mathbb{N})\text{ when }|z|=1.$$

Here $\mathbb{Z}_0^-=\{0,-1,-2,\ldots\}$, $\mathbb{N}$ is the set of positive integers, and $\mathbb{C}$ is the set of complex numbers (see [2]).

Letting $z=1$ in (1.1), we have the Hurwitz zeta function

$$\zeta(s,a)=\sum_{n=0}^{\infty}\frac{1}{(a+n)^s}. \tag{1.2}$$

Letting $a=1$ in (1.2), we obtain the Riemann zeta function

$$\zeta_{\mathbb{Q}}(s)=\sum_{n=1}^{\infty}\frac{1}{n^s}. \tag{1.3}$$

Let $P_1(x)=B_1(x-[x])=x-[x]-1/2$ be the first periodized Bernoulli polynomials, and $\{x\}=x-[x]$ be the fractional part of x. Recently, by using Euler-MacLaurin summation formula

$$\sum_{\alpha<n\le\beta}\Phi(n)=\int_\alpha^\beta\Phi(x)dx+\int_\alpha^\beta\Phi'(x)P_1(x)dx+P_1(\alpha)\Phi(\alpha)-P_1(\beta)\Phi(\beta), \tag{1.4}$$

Coffey [1, p. 81] gave an integral representation of the Hurwitz-Lerch zeta function $\Phi(z,s,a)$ (1.1) as follows.

2000 *Mathematics Subject Classification.* 11F03,11M35,33E05.

Key words and phrases. Integral representation, Elliptic function.

Proposition 1.1 (Coffey [1])**.**

$$\Phi(z,s,a)=\frac{1}{a^s}+\frac{z}{2(a+1)^s}+\int_1^\infty\frac{z^x}{(x+a)^s}dx+\int_1^\infty\left[\frac{z^x\ln z}{(x+a)^s}-\frac{sz^x}{(x+a)^{s+1}}\right]P_1(x)dx. \tag{1.5}$$

Suppose W is a lattice in the complex plane, ω_1 and ω_2 are two generators of W, so that W consists of the points $w=n\omega_1+m\omega_2$, where n and m are integers. In his historical book "Elliptic functions according to Eisenstein and Kronecker" [3], generalizing the Hurwitz zeta functions (1.2) above, A. Weil [3, p. 14] introduced the following elliptic function:

$$E_k(a,W)=\sum_{w\in W}\frac{1}{(a+w)^k}=\sum_{(n,m)\in\mathbb{Z}^2}\frac{1}{(a+n\omega_1+m\omega_2)^k} \tag{1.6}$$

for $a\notin W$, which is also a generalization of the homogeneous Eisenstein series defined by

$$G_k(W)=\sum_{0\neq w\in W}\frac{1}{w^k}=\sum_{(n,m)\in\mathbb{Z}^2\backslash(0,0)}\frac{1}{(n\omega_1+m\omega_2)^k}. \tag{1.7}$$

As pointed out by Weil [3, p. 14], the series (1.6) are absolutely convergent for $k\geq 3$. If $k=1$ and $k=2$, then Eisenstein makes use of a summatory process which we shall call Eisenstein summation., that is,

$$\sum_{(n,m)\in\mathbb{Z}^2}\frac{1}{(a+n\omega_1+m\omega_2)^k}=\lim_{M\to\infty}\sum_{m=-M}^{m=M}\left(\lim_{N\to\infty}\sum_{n=-N}^{N}\frac{1}{(a+n\omega_1+m\omega_2)^k}\right). \tag{1.8}$$

In order to give an analogue of the above Coffey's result for Weil's elliptic functions (1.6), in this paper, we shall generalize Euler-MacLaurin summation formula (1.4) to a two dimensional case.

Proposition 1.2 (Two dimensional summation formula)**.** *Suppose* $\Phi(x,y)=f(a+x\omega_1+y\omega_2)\in C^2([\alpha_1,\beta_1]\times[\alpha_2,\beta_2])$*, we have*

$$\sum_{\alpha_2<m\leq\beta_2}\sum_{\alpha_1<n\leq\beta_1}\Phi(n,m)=I_1+I_2+I_3+I_4, \tag{1.9}$$

where

$$I_1=\iint_{[\alpha_1,\beta_1]\times[\alpha_2,\beta_2]}\left[\Phi(x,y)+\frac{\partial\Phi(x,y)}{\partial x}P_1(x)+\frac{\partial\Phi(x,y)}{\partial y}P_1(y)+\frac{\partial^2\Phi(x,y)}{\partial x\partial y}P_1(x)P_1(y)\right]dxdy, \tag{1.10}$$

$$I_2=\int_{\alpha_2}^{\beta_2}\left[\Phi(\alpha_1,y)P_1(\alpha_1)-\Phi(\beta_1,y)P_1(\beta_1)+\frac{\partial\Phi(\alpha_1,y)}{\partial y}P_1(y)P_1(\alpha_1)-\frac{\partial\Phi(\beta_1,y)}{\partial y}P_1(y)P_1(\beta_1)\right]dy, \tag{1.11}$$

$$
\begin{aligned}
I_3 = \int_{\alpha_1}^{\beta_1} \Bigg[& \Phi(x,\alpha_2)P_1(\alpha_2) - \Phi(x,\beta_2)P_1(\beta_2) \\
& + \frac{\partial \Phi(x,\alpha_2)}{\partial x} P_1(x)P_1(\alpha_2) - \frac{\partial \Phi(x,\beta_2)}{\partial x} P_1(x)P_1(\beta_2) \Bigg] dx,
\end{aligned}
\tag{1.12}
$$

and

$$
\begin{aligned}
I_4 = {} & P_1(\alpha_2)P_1(\alpha_1)\Phi(\alpha_1,\alpha_2) - P_1(\alpha_2)P_1(\beta_1)\Phi(\beta_1,\alpha_2) \\
& - P_1(\beta_2)P_1(\alpha_1)\Phi(\alpha_1,\beta_2) + P_1(\beta_2)P_1(\beta_1)\Phi(\beta_1,\beta_2).
\end{aligned}
\tag{1.13}
$$



Remark 1.3. From the above graph, we see that I_1, the first term in the right side of the above formula, is an integral inside the square $[\alpha_1,\beta_1] \times [\alpha_2,\beta_2]$. The second term I_2 and the third term I_3 are the integral along the boundary lines $[\alpha_2,\beta_2]$ and $[\alpha_1,\beta_1]$, respectively. The fourth term I_4 is a sum at the four corner points $(\alpha_1,\alpha_2),(\beta_1,\alpha_2),(\alpha_1,\beta_2),(\beta_1,\beta_2)$.

Thus I_1 is corresponding to $\int_\alpha^\beta \Phi(x)dx + \int_\alpha^\beta \Phi'(x)P_1(x)dx$ (an integral inside the interval $[\alpha,\beta]$) in the Euler-MacLaurin summation formula (see (1.4) above), while the term $I_2+I_3+I_4$ is corresponding to the term $P_1(\alpha)\Phi(\alpha)-P_1(\beta)\Phi(\beta)$ (a sum at the boundary points of the interval $[\alpha,\beta]$) in the Euler-MacLaurin summation formula (1.4).

The following is an integral representation of Weil's elliptic function (1.6) which is an analogue of Proposition 1.1 on the integral representation of the Hurwitz-Lerch zeta function.

Corollary 1.4. *Suppose* $(x_0,y_0)\in\mathbb{R}^2$ *is the point such that* $x_0w_1+y_0w_2=-a$*, for arbitrary small* $\epsilon>0$*, we have*

$$
E_k(a,W) = \sum_{(n,m)\in\mathbb{Z}^2} \frac{1}{(a+n\omega_1+m\omega_2)^k} = J_1+J_2+J_3,
\tag{1.14}
$$

where

$$\begin{aligned} J_1 = \int_{-\infty}^{\infty} \bigg[& \frac{1}{(a+xw_1+(y_0+\epsilon)w_2)^k} P_1(y_0+\epsilon) \\ & - \frac{1}{(a+xw_1+(y_0-\epsilon)w_2)^k} P_1(y_0-\epsilon) \\ & + k\frac{w_1}{(a+xw_1+(y_0-\epsilon)w_2)^{k+1}} P_1(x)P_1(y_0-\epsilon) \\ & - k\frac{w_1}{(a+xw_1+(y_0+\epsilon)w_2)^{k+1}} P_1(x)P_1(y_0+\epsilon) \bigg] dx, \end{aligned} \tag{1.15}$$

$$\begin{aligned} J_2 = \iint_{[-\infty,\infty]\times[y_0+\epsilon,\infty]} \bigg[& \frac{1}{(a+xw_1+yw_2)^k} \\ & - k\frac{w_1}{(a+xw_1+yw_2)^{k+1}} P_1(x) \\ & - k\frac{w_2}{(a+xw_1+yw_2)^{k+1}} P_1(y) \\ & + k(k+1)\frac{w_1w_2}{(a+xw_1+yw_2)^{k+2}} P_1(x)P_1(y) \bigg] dxdy, \end{aligned} \tag{1.16}$$

and

$$\begin{aligned} J_3 = \iint_{[-\infty,\infty]\times[-\infty,y_0-\epsilon]} \bigg[& \frac{1}{(a+xw_1+yw_2)^k} \\ & - k\frac{w_1}{(a+xw_1+yw_2)^{k+1}} P_1(x) \\ & - k\frac{w_2}{(a+xw_1+yw_2)^{k+1}} P_1(y) \\ & + k(k+1)\frac{w_1w_2}{(a+xw_1+yw_2)^{k+2}} P_1(x)P_1(y) \bigg] dxdy. \end{aligned} \tag{1.17}$$



Remark 1.5. If $k=1$ and $k=2$, then the integral $\int_{-\infty}^{\infty}$ should be understood as

$$\int_{-\infty}^{\infty} = \lim_{N\to\infty}\int_{-N}^{N}.$$

In Sections 2 and 3, we shall prove Proposition 1.2 and Corollary 1.4, respectively.

2. PROOF OF PROPOSITION 1.2

The following Euler-MacLaurin summation formula will be used many times in our proof.

$$\sum_{\alpha<n\le\beta}\Phi(n)=\int_\alpha^\beta\Phi(x)dx+\int_\alpha^\beta\Phi'(x)P_1(x)dx+P_1(\alpha)\Phi(\alpha)-P_1(\beta)\Phi(\beta). \tag{2.1}$$

By taking $\Phi(y)=\sum_{\alpha_1<n\le\beta_1}f(a+nw_1+yw_2)$ in (2.1), we have

$$\begin{aligned}\sum_{\alpha_2<m\le\beta_2}\left(\sum_{\alpha_1<n\le\beta_1}f(a+nw_1+mw_2)\right)&=\int_{\alpha_2}^{\beta_2}\left(\sum_{\alpha_1<n\le\beta_1}f(a+nw_1+yw_2)\right)dy\\&+\int_{\alpha_2}^{\beta_2}\left(\sum_{\alpha_1<n\le\beta_1}\frac{\partial f(a+nw_1+yw_2)}{\partial y}\right)P_1(y)dy\\&+P_1(\alpha_2)\sum_{\alpha_1<n\le\beta_1}f(a+nw_1+\alpha_2w_2)\\&-P_1(\beta_2)\sum_{\alpha_1<n\le\beta_1}f(a+nw_1+\beta_2w_2).\end{aligned}\tag{2.2}$$

By taking $\Phi(x)=f(a+xw_1+yw_2)$ in (2.1), we have

$$\begin{aligned}\sum_{\alpha_1<n\le\beta_1}f(a+nw_1+mw_2)&=\int_{\alpha_1}^{\beta_1}f(a+xw_1+yw_2)dx\\&+\int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+yw_2)}{\partial x}P_1(x)dx\\&+P_1(\alpha_1)f(a+\alpha_1w_1+yw_2)\\&-P_1(\beta_1)f(a+\beta_1w_1+yw_2).\end{aligned}\tag{2.3}$$

By taking $\Phi(x)=\frac{\partial f(a+xw_1+yw_2)}{\partial y}$ in (2.1), we have

$$\begin{aligned}\sum_{\alpha_1<n\le\beta_1}\frac{\partial f(a+nw_1+yw_2)}{\partial y}&=\int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+yw_2)}{\partial y}dx\\&+\int_{\alpha_1}^{\beta_1}\frac{\partial^2 f(a+xw_1+yw_2)}{\partial x\partial y}P_1(x)dx\\&+P_1(\alpha_1)\frac{\partial f(a+\alpha_1w_1+yw_2)}{\partial y}\\&-P_1(\beta_1)\frac{\partial f(a+\beta_1w_1+yw_2)}{\partial y}.\end{aligned}\tag{2.4}$$

By taking $\Phi(x)=f(a+xw_1+\alpha_2w_2)$ in (2.1), we have

$$\begin{aligned}\sum_{\alpha_1<n\leq\beta_1} f(a+nw_1+\alpha_2w_2) = &\int_{\alpha_1}^{\beta_1} f(a+xw_1+\alpha_2w_2)dx \\ &+\int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+\alpha_2w_2)}{\partial x}P_1(x)dx \\ &+P_1(\alpha_1)f(a+\alpha_1w_1+\alpha_2w_2) \\ &-P_1(\beta_1)f(a+\beta_1w_1+\alpha_2w_2).\end{aligned}\tag{2.5}$$

By taking $\Phi(x)=f(a+xw_1+\beta_2w_2)$ in (2.1), we have

$$\begin{aligned}\sum_{\alpha_1<n\leq\beta_1} f(a+nw_1+\beta_2w_2) = &\int_{\alpha_1}^{\beta_1} f(a+xw_1+\beta_2w_2)dx \\ &+\int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+\beta_2w_2)}{\partial x}P_1(x)dx \\ &+P_1(\alpha_1)f(a+\alpha_1w_1+\beta_2w_2) \\ &-P_1(\beta_1)f(a+\beta_1w_1+\beta_2w_2).\end{aligned}\tag{2.6}$$

Substituting (2.3), (2.4), (2.5), (2.6) into (2.2), we have

$$
\begin{aligned}
\sum_{\alpha_2<m\leq\beta_2}\sum_{\alpha_1<n\leq\beta_1} f(a+nw_1+mw_2) &= \sum_{\alpha_2<m\leq\beta_2}\left(\sum_{\alpha_1<n\leq\beta_1} f(a+nw_1+mw_2)\right)\\
&= \int_{\alpha_2}^{\beta_2}\Bigg[\int_{\alpha_1}^{\beta_1} f(a+xw_1+yw_2)dx\\
&\quad + \int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+yw_2)}{\partial x}P_1(x)dx\\
&\quad + P_1(\alpha_1)f(a+\alpha_1w_1+yw_2)\\
&\quad - P_1(\beta_1)f(a+\beta_1w_1+yw_2)\Bigg]dy\\
&+ \int_{\alpha_2}^{\beta_2}\Bigg[\int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+yw_2)}{\partial y}dx\\
&\quad + \int_{\alpha_1}^{\beta_1}\frac{\partial^2 f(a+xw_1+yw_2)}{\partial x\partial y}P_1(x)dx\\
&\quad + P_1(\alpha_1)\frac{\partial f(a+\alpha_1w_1+yw_2)}{\partial y}\\
&\quad - P_1(\beta_1)\frac{\partial f(a+\beta_1w_1+yw_2)}{\partial y}\Bigg]P_1(y)dy\\
&+ P_1(\alpha_2)\Bigg[\int_{\alpha_1}^{\beta_1} f(a+xw_1+\alpha_2w_2)dx\\
&\quad + \int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+\alpha_2w_2)}{\partial x}P_1(x)dx\\
&\quad + P_1(\alpha_1)f(a+\alpha_1w_1+\alpha_2w_2)\\
&\quad - P_1(\beta_1)f(a+\beta_1w_1+\alpha_2w_2)\Bigg]\\
&- P_1(\beta_2)\Bigg[\int_{\alpha_1}^{\beta_1} f(a+xw_1+\beta_2w_2)dx\\
&\quad + \int_{\alpha_1}^{\beta_1}\frac{\partial f(a+xw_1+\beta_2w_2)}{\partial x}P_1(x)dx\\
&\quad + P_1(\alpha_1)f(a+\alpha_1w_1+\beta_2w_2)\\
&\quad - P_1(\beta_1)f(a+\beta_1w_1+\beta_2w_2)\Bigg]
\end{aligned}
$$

This equals to

$$\begin{aligned}
&\int_{\alpha_2}^{\beta_2}\int_{\alpha_1}^{\beta_1} f(a+xw_1+yw_2)dxdy\\
&\quad+\int_{\alpha_2}^{\beta_2}\int_{\alpha_1}^{\beta_1} \frac{\partial f(a+xw_1+yw_2)}{\partial x}P_1(x)dxdy\\
&\quad+P_1(\alpha_1)\int_{\alpha_2}^{\beta_2} f(a+\alpha_1w_1+yw_2)dy\\
&\quad-P_1(\beta_1)\int_{\alpha_2}^{\beta_2} f(a+\beta_1w_1+yw_2)dy\\
&\quad+\int_{\alpha_2}^{\beta_2}\int_{\alpha_1}^{\beta_1} \frac{\partial f(a+xw_1+yw_2)}{\partial y}P_1(y)dxdy\\
&\quad+\int_{\alpha_2}^{\beta_2}\int_{\alpha_1}^{\beta_1} \frac{\partial^2 f(a+xw_1+yw_2)}{\partial x\partial y}P_1(x)P_1(y)dxdy\\
&\quad+P_1(\alpha_1)\int_{\alpha_2}^{\beta_2} \frac{\partial f(a+\alpha_1w_1+yw_2)}{\partial y}P_1(y)dy\\
&\quad-P_1(\beta_1)\int_{\alpha_2}^{\beta_2} \frac{\partial f(a+\beta_1w_1+yw_2)}{\partial y}P_1(y)dy\\
&\quad+P_1(\alpha_2)\int_{\alpha_1}^{\beta_1} f(a+xw_1+\alpha_2w_2)dx\\
&\quad+P_1(\alpha_2)\int_{\alpha_1}^{\beta_1} \frac{\partial f(a+xw_1+\alpha_2w_2)}{\partial x}P_1(x)dx\\
&\quad+P_1(\alpha_2)P_1(\alpha_1)f(a+\alpha_1w_1+\alpha_2w_2)\\
&\quad-P_1(\alpha_2)P_1(\beta_1)f(a+\beta_1w_1+\alpha_2w_2)\\
&\quad-P_1(\beta_2)\int_{\alpha_1}^{\beta_1} f(a+xw_1+\beta_2w_2)dx\\
&\quad-P_1(\beta_2)\int_{\alpha_1}^{\beta_1} \frac{\partial f(a+xw_1+\beta_2w_2)}{\partial x}P_1(x)dx\\
&\quad-P_1(\beta_2)P_1(\alpha_1)f(a+\alpha_1w_1+\beta_2w_2)\\
&\quad+P_1(\beta_2)P_1(\beta_1)f(a+\beta_1w_1+\beta_2w_2).
\end{aligned}$$

Finally, we have

$$
\begin{aligned}
\sum_{\alpha_2<m\leq\beta_2}\sum_{\alpha_1<n\leq\beta_1} f(a+nw_1+mw_2) = {} & \int_{\alpha_2}^{\beta_2}\int_{\alpha_1}^{\beta_1}\Bigg[f(a+xw_1+yw_2) \\
& + \frac{\partial f(a+xw_1+yw_2)}{\partial x}P_1(x) \\
& + \frac{\partial f(a+xw_1+yw_2)}{\partial y}P_1(y) \\
& + \frac{\partial^2 f(a+xw_1+yw_2)}{\partial x\partial y}P_1(x)P_1(y)\Bigg]dxdy \\
& + \int_{\alpha_2}^{\beta_2}\Bigg[f(a+\alpha_1w_1+yw_2)P_1(\alpha_1) \\
& - f(a+\beta_1w_1+yw_2)P_1(\beta_1) \\
& + \frac{\partial f(a+\alpha_1w_1+yw_2)}{\partial y}P_1(y)P_1(\alpha_1) \\
& - \frac{\partial f(a+\beta_1w_1+yw_2)}{\partial y}P_1(y)P_1(\beta_1)\Bigg]dy \\
& + \int_{\alpha_1}^{\beta_1}\Bigg[f(a+xw_1+\alpha_2w_2)P_1(\alpha_2) \\
& - f(a+xw_1+\beta_2w_2)P_1(\beta_2) \\
& + \frac{\partial f(a+xw_1+\alpha_2w_2)}{\partial x}P_1(x)P_1(\alpha_2) \\
& - \frac{\partial f(a+xw_1+\beta_2w_2)}{\partial x}P_1(x)P_1(\beta_2)\Bigg]dx \\
& + P_1(\alpha_2)P_1(\alpha_1)f(a+\alpha_1w_1+\alpha_2w_2) \\
& - P_1(\alpha_2)P_1(\beta_1)f(a+\beta_1w_1+\alpha_2w_2) \\
& - P_1(\beta_2)P_1(\alpha_1)f(a+\alpha_1w_1+\beta_2w_2) \\
& + P_1(\beta_2)P_1(\beta_1)f(a+\beta_1w_1+\beta_2w_2).
\end{aligned} \tag{2.7}
$$

Notice that $\Phi(x,y)=f(a+x\omega_1+y\omega_2)$, we get our result.

3. PROOF OF COROLLARY 1.4

In this section, let

$$f(a,x,y)=\frac{1}{(a+xw_1+yw_2)^k}. \tag{3.1}$$

We have

$$\frac{\partial f}{\partial x}(x,y)=-k\frac{w_1}{(a+xw_1+yw_2)^{k+1}}, \tag{3.2}$$

$$\frac{\partial^2 f}{\partial x\partial y}(x,y)=k(k+1)\frac{w_1w_2}{(a+xw_1+yw_2)^{k+2}}, \tag{3.3}$$

$$\frac{\partial f}{\partial y}(x,y) = -k\frac{w_2}{(a+xw_1+yw_2)^{k+1}}, \tag{3.4}$$

$$\frac{\partial f}{\partial x}(N,y) = -k\frac{w_1}{(a+Nw_1+yw_2)^{k+1}} \tag{3.5}$$

and

$$\frac{\partial f}{\partial y}(x,N) = -k\frac{w_2}{(a+xw_1+Nw_2)^{k+1}}. \tag{3.6}$$

We have

$$\begin{aligned} |f(a,N,y)| &= \frac{1}{|a+Nw_1+yw_2|^k} \to 0, \\ \left|\frac{\partial f}{\partial x}(N,y)\right| &= k\frac{|w_1|}{|a+Nw_1+yw_2|^{k+1}} \to 0 \quad \text{as } N\to\infty, \end{aligned} \tag{3.7}$$

uniformly for $y \in [\alpha_2, \beta_2]$, and

$$\begin{aligned} |f(a,x,N)| &= \frac{1}{|a+xw_1+Nw_2|^k} \to 0, \\ \left|\frac{\partial f}{\partial y}(x,N)\right| &= k\frac{|w_2|}{|a+xw_1+Nw_2|^{k+1}} \to 0 \quad \text{as } N\to\infty, \end{aligned} \tag{3.8}$$

uniformly for $x \in [\alpha_1, \beta_1]$.

Thus, by taking $\alpha_1 = -N$, $\beta_1 = N$, $\alpha_2 = y_0+\epsilon$, $\beta_2 = N$ and $f(a,x,y) = \frac{1}{(a+xw_1+yw_2)^k}$ in (2.7), then letting $N \to \infty$, we have

$$\begin{aligned} \sum_{y_0+\epsilon<m<\infty}\ \sum_{-\infty<n<\infty} \frac{1}{(a+nw_1+mw_2)^k} = \int_{y_0+\epsilon}^{\infty}\int_{-\infty}^{\infty} \Bigg[& \frac{1}{(a+xw_1+yw_2)^k} \\ & - k\frac{w_1}{(a+xw_1+yw_2)^{k+1}}P_1(x) \\ & - k\frac{w_2}{(a+xw_1+yw_2)^{k+1}}P_1(y) \\ & + k(k+1)\frac{w_1w_2}{(a+xw_1+yw_2)^{k+2}}P_1(x)P_1(y)\Bigg]dxdy \\ + \int_{-\infty}^{\infty}\Bigg[& \frac{1}{(a+xw_1+(y_0+\epsilon)w_2)^k}P_1(y_0+\epsilon) \\ & - k\frac{w_1}{(a+xw_1+(y_0+\epsilon)w_2)^{k+1}}P_1(x)P_1(y_0+\epsilon)\Bigg]dx. \end{aligned} \tag{3.9}$$

Similarly, we have
(3.10)

$$\begin{aligned}\sum_{-\infty<m\leq y_0-\epsilon}\ \sum_{-\infty<n<\infty}\frac{1}{(a+nw_1+mw_2)^k}=&\int_{-\infty}^{y_0-\epsilon}\int_{-\infty}^{\infty}\bigg[\frac{1}{(a+xw_1+yw_2)^k}\\&-k\frac{w_1}{(a+xw_1+yw_2)^{k+1}}P_1(x)\\&-k\frac{w_2}{(a+xw_1+yw_2)^{k+1}}P_1(y)\\&+k(k+1)\frac{w_1w_2}{(a+xw_1+yw_2)^{k+2}}P_1(x)P_1(y)\bigg]dxdy\\&-\int_{-\infty}^{\infty}\bigg[\frac{1}{(a+xw_1+(y_0-\epsilon)w_2)^k}P_1(y_0-\epsilon)\\&+k\frac{w_1}{(a+xw_1+(y_0-\epsilon)w_2)^{k+1}}P_1(x)P_1(y_0-\epsilon)\bigg]dx.\end{aligned}$$

Since $a\notin W$, for arbitrary small $\epsilon>0$, we have
(3.11)

$$\begin{aligned}\sum_{(n,m)\in\mathbb{Z}^2}\frac{1}{(a+n\omega_1+m\omega_2)^k}=&\sum_{y_0+\epsilon<m<\infty}\ \sum_{-\infty<n<\infty}\frac{1}{(a+nw_1+mw_2)^k}\\&+\sum_{-\infty<m\leq y_0-\epsilon}\ \sum_{-\infty<n<\infty}\frac{1}{(a+nw_1+mw_2)^k},\end{aligned}$$

from (3.9) and (3.10), we get our result.

REFERENCES

[1] M.W. Coffey, *Integral representations of functions and Addison-type series for mathematical constants*, J. Number Theory **157** (2015), 79–98.

[2] H. M. Srivastava and J. Choi, Zeta and q-Zeta Functions and Associated Series and Integrals, Elsevier Science Publishers, Amsterdam, London and New York, 2012.

[3] A. Weil, Elliptic functions according to Eisenstein and Kronecker, Reprint of the 1976 original, Classics in Mathematics, Springer-Verlag, Berlin, 1999.

DEPARTMENT OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY, GUANGZHOU, GUANGDONG 510640, CHINA
E-mail address: `hus04@mails.tsinghua.edu.cn`

CENTER FOR GENERAL EDUCATION, KYUNGNAM UNIVERSITY, 7(WORYEONG-DONG) KYUNGNAMDAEHAK-RO, MASANHAPPO-GU, CHANGWON-SI, GYEONGSANGNAM-DO 631-701, REPUBLIC OF KOREA
E-mail address: `mskim@kyungnam.ac.kr`